

5

**THERAPEUTIC DELIVERY COMPOSITIONS
AND METHODS OF USE THEREOF**

10 **Cross-Reference to Related Applications**

15 This application is a continuation-in-part of co-
pending U.S. Application Serial No. 09/457,771, filed
December 9, 1999, which is a continuation of U.S. Application
Serial No. 09/104,088, filed June 24, 1998, now abandoned,
which is a continuation-in-part of U.S. Application Serial No.
08/926,297, filed September 5, 1997, now abandoned, which is
a continuation of U.S. Application Serial No. 08/725,842, filed
September 30, 1996, now abandoned, which is a continuation of
U.S. Application Serial No. 08/138,271, filed October 15, 1993,
20 now abandoned.

25 This application also claims priority under 35
U.S.C. § 120 to co-pending U.S. Application Serial No.
09/368,855, filed August 5, 1999, which is a continuation of
U.S. Application Serial No. 08/889,342 filed July 8, 1997, now
United States Patent No. 5,990,241, which is a continuation of
U.S. Application Serial No. 08/657,161, filed June 3, 1996, now
United States Patent No. 5,691,387, which is a division of U.S.
Application Serial No. 08/087,136 filed July 2, 1993, now
30 United States Patent No. Re. 36,665, which is a continuation of
U.S. Application Serial No. 07/847,874 filed March 13, 1992,
now abandoned, which is a continuation-in-part of United States
Application Serial No. 07/673,289, filed March 19, 1991, now
abandoned.

35

Technical Field

5 The present invention relates to therapeutic delivery compounds and to compositions comprising therapeutic delivery compounds that kill or suppress the growth of bacteria, viruses, fungi and protozoa, and methods of use thereof. The compounds, compositions and methods are effective for the delivery of drugs and other compounds to the interior of cells and for controlling intracellular organisms.

10 Background of the Invention

Many new and potentially useful technologies are being developed which may form the basis of future medical cures and therapies. Examples of such technologies include, gene replacement, antisense gene therapy, triplex gene therapy and ribozyme-based therapy. However, to be successful, these technologies require effective means for the delivery of the therapeutic agent across cellular, nuclear and microorganismal membranes.

20 The recent advent of technology, and advances in our ability to understand the structure and function of many genes makes it possible to selectively turn off or modify the activity of a given gene. Alteration of gene activity can be accomplished many ways. For example, oligonucleotides that are complementary to certain gene messages or viral sequences, known as "antisense" compounds, have been shown to have an inhibitory effect against viruses. By creating an antisense compound that hybridizes with the targeted RNA message of cells or viruses the translation of the message into protein can be interrupted or prevented. In this fashion gene activity can be modulated.

30 The ability to deactivate specific genes provides great therapeutic benefits. For example, it is theoretically possible to fight viral diseases with antisense RNA and DNA molecules that seek out and destroy viral gene products. In tissue culture, antisense oligonucleotides have inhibited

35

infections by herpes-viruses, influenza viruses and the human immunodeficiency virus that causes AIDS. It may also be possible to target antisense oligonucleotides against mutated oncogenes. Antisense technology also holds the potential for regulating growth and development. However, in order for the gene therapy to work, antisense therapeutic compounds must be delivered across cellular plasma membranes to the cytosol.

Gene activity is also modified using sense DNA in a technique known as gene therapy. Defective genes are replaced or supplemented by the administration of "good" or normal genes that are not subject to the defect. The administered normal genes which insert into a chromosome, or may be present in extracellular DNA, produce normal RNA, which in turn leads to normal gene product. In this fashion gene defects and deficiencies in the production of gene product may be corrected. Still further gene therapy has the potential to augment the normal genetic complement of a cell. For example, it has been proposed that one way to combat HIV is to introduce into an infected person's T cells a gene that makes the cells resistant to HIV infection. This form of gene therapy is sometimes called "intracellular immunization." Genetic material such as polynucleotides may be administered to a mammal to elicit an immune response against the gene product of the administered nucleic acid sequence. Such gene vaccines elicit an immune response in the following manner. First, the nucleic acid sequence is administered to a human or animal. Next, the administered sequence is expressed to form gene product within the human or animal. The gene product inside the human or animal is recognized as foreign material and the immune system of the human or animal mounts an immunological response against the gene product. However, this approach currently is not feasible due to a lack of effective gene delivery systems that facilitate the delivery of genetic material across both cellular and nuclear membranes.

Finally, gene therapy may be used as a method of delivering drugs *in vivo*. For example, if genes that code for therapeutic compounds can be delivered to endothelial cells, the gene products would have facilitated access to the blood stream. Currently, genes are delivered to cells *ex vivo* and then reintroduced to the animal.

Retroviral vectors can be used to deliver genes *ex vivo* to isolated cells, which are then infused back into the patient. However, retroviral vectors have some drawbacks, such as being able to deliver genes only to dividing cells, random integration of the gene to be delivered, potentially causing unwanted genetic alterations, and possibly reverting back to an infectious wild-type retroviral form. Another drawback of antisense gene therapy is that it is effective at the messenger RNA level, which means that antisense oligonucleotides must be introduced in a quantity to interact with all or a substantial number of the mRNA in the cytosol, and that such treatment is only effective during active synthesis of mRNA. Further, the oligonucleotides must be maintained at this high quantity level throughout mRNA synthesis to be effective over time.

Newly developed "triplex DNA" technology represents an improvement in gene regulation. Triplex DNA technology utilizes oligonucleotides and compounds that specifically bind to particular regions of duplex DNA, thereby inactivating the targeted gene. An advantage of triplex DNA technology is that only a single copy of the oligonucleotide or compound is required to alter gene expression because the binding is at the DNA level, not the mRNA level. A drawback of triplex DNA technology, however, is that the oligonucleotide or compound must pass through not only the cellular membrane, but also the microbial membrane in the case of treating microbial infections, or the nuclear membrane in the case of altering eukaryotic gene function or expression of foreign DNA integrated into chromosomal DNA.

Another emerging technology relates to the therapeutic use of ribozymes for the treatment of genetic disorders. Ribozymes are catalytic RNA molecules that consist of a hybridizing region and an enzymatic region. Ribozymes may in the future be engineered so as to specifically bind to a targeted region of nucleic acid sequence and cut or otherwise enzymatically modify the sequence so as to alter its expression or translation into gene product.

There is a great need, therefore, for improved delivery systems for genetic material such as genes, polynucleotides, and antisense oligonucleotides that can be used in gene therapy. More specifically, there is a need for non-toxic compositions having surfactant properties that can facilitate the transport of genetic compounds and other drugs and therapeutic compounds across cellular membranes.

There is a particularly urgent need for an effective treatment for Acquired Immune Deficiency Syndrome, or AIDS, a disease thought to be caused by a human retrovirus, the Human T Lymphotropic Virus III (HTLV-III) which is also called human immunodeficiency virus or HIV. Like other retroviruses, HIV has ribonucleic acid, or RNA, as its genetic material. When the virus enters the host cell, a viral enzyme called reverse transcriptase exploits the viral RNA as a template to assemble a corresponding molecule of DNA. The DNA travels through the cell nucleus and inserts itself among the host chromosomes, where it provides the basis for viral replication.

In the case of HIV, the host cell is often a T4 lymphocyte, a white blood cell that has a central and regulatory role in the immune system. Once it is inside a T4 cell, the virus may remain latent until the lymphocyte is immunologically stimulated by a secondary infection. Then the virus reproducing itself rapidly killing or rendering ineffective the host cell. The resulting depletion of the T4 cells, and loss of activity leaves the patient vulnerable to "opportunistic" infections by an agent that

would not normally harm a healthy person. The virus damages the host by many other mechanisms as well.


Many therapies against AIDS infection that are currently being investigated. Several of these therapies under investigation are based on interrupting the reverse transcriptase as it assembles the viral DNA destined to become the virus. The drugs used for this purpose are chemical analogs of the nucleic acids that form the subunits of DNA. When the analog is supplied to an infected cell, reverse transcriptase will incorporate it into a growing DNA chain. Because the analog lacks the correct attachment point for the next subunit, however, the chain is terminated. The truncated DNA cannot integrate itself into the host chromosomes or provide the basis for viral replication, and so the spread of the infection is halted. One of the compounds that is thought to act by mimicking a nucleotide is azidothymidine, or AZT. However, AZT is known to have serious side effects and its efficacy in mitigating the AIDS disease has been questioned. The efficacy of AZT and other antiviral and antimicrobial drugs could be increased if improved means and methods for delivering therapeutic agents to the site of infection were available.

Summary of the Invention

The present invention includes a method of delivering therapeutic drugs to a human or animal for treating disease states such as, but not limited to, bacterial infection and infections caused by HIV and other DNA and RNA viruses. The present invention relates particularly to compositions and methods for treating infectious diseases and genetic disorders through gene therapy and intracellular delivery of antisense oligonucleotides or other nucleic acid sequences.

The present invention comprises a therapeutic delivery composition effective for treating a disease state comprising an administerable admixture of an effective amount of a therapeutic compound capable of altering nucleic acid

sequence function and an effective amount of a surface active nonionic block copolymer having the following general formula:



5


wherein a is an integer such that the hydrophobe represented by (C₃H₆O) has a molecular weight of approximately 750 and approximately 15,000, preferably between approximately 2250 and approximately 15,000, more preferably between approximately 3250 and approximately 15,000, and b is an integer such that the hydrophile portion represented by (C₂H₄O) constitutes approximately 1% to approximately 50% by weight of the compound, preferably approximately 5% to approximately 20%.

10

15

Another aspect of the present invention comprises the surface active nonionic block copolymer of the present invention also comprises a reverse triblock copolymer of polyoxypropylene/polyoxyethylene having the formula:

20




25

wherein "b" represents a number such that the molecular weight of the hydrophobe (C₃H₆O)_b is between approximately 2,000 and 10,000, and "a" represents a number such that the percentage of hydrophile (C₂H₄O)_a is between approximately 2% and 30%.

30

Another embodiment of the surface active nonionic block copolymer of the present invention, comprises the reverse triblock copolymer of polyoxypropylene/polyoxyethylene having the formula:




35

wherein "b" represents a number such that the molecular weight of the hydrophobe (C₃H₆O)_b is between approximately


2,500 and 8,500, and "a" represents a number such that the percentage of hydrophile $(C_2H_4O)_a$ is between approximately 7% and 23%.

5 Yet another embodiment of the present invention, comprises the reverse triblock copolymer of polyoxypropylene/polyoxyethylene having the formula:



10 wherein "b" represents a number such that the molecular weight of the hydrophobe $(C_3H_6O)_b$ is between approximately 3,000 and 7,000, and "a" represents a number such that the percentage of hydrophile $(C_2H_4O)_a$ is between approximately 8% and 20%.

15 A preferred embodiment of the present invention, comprises the reverse triblock copolymer polyoxypropylene/polyoxyethylene having the formula:




20 wherein "b" represents a number such that the molecular weight of the hydrophobe $(C_3H_6O)_b$ is between approximately 5,000 and 6,000, and "a" represents a number such that the percentage of hydrophile $(C_2H_4O)_a$ is between approximately 14% and 16%.

25 In general, "a" represents a number such that the percentage of hydrophile $(C_2H_4O)_a$ is between approximately 5% and 30%, with a preferable range of between approximately 7% and 23% and a most preferable range of between 8% and 20%.

30 In another aspect, the present invention comprises a therapeutic delivery composition effective for treating a disease state comprising an administerable admixture of an effective amount of a therapeutic compound capable of altering
35 nucleic acid sequence function and an effective amount of a

biologically-active copolymer comprising a copolymer of polyoxyethylene (POE) which is hydrophilic and polyoxypropylene (POP) which is hydrophobic. The block copolymer is built on a tetrafunctional ethylenediamine initiator. In the preferred embodiment of the biologically-active copolymers of the present invention, the block copolymers that comprise the biologically-active copolymers of the present invention have the following general formulas:



wherein:


the mean aggregate molecular weight of the hydrophobe portion of the octablock copolymer consisting of polyoxypropylene (C₃H₆O)_b (POP) is between approximately 5000 and 7000 daltons;

a is a number such that the hydrophile portion represented by polyoxyethylene (C₂H₄O)_a (POE) constitutes between approximately 10% and 40% of the total molecular weight of the compound; and

b is a number such that the polyoxypropylene (C₃H₆O)_b (POP) portion of the total molecular weight of the octablock copolymer constitutes between approximately 60% and 90% of the compound.

In another aspect of the present invention, the biologically-active copolymer comprises a polymer of hydrophilic polyoxyethylene (POE) built on an ethylene diamine initiator. Polymers of hydrophobic polyoxypropylene (POP) are then added to the blocks of hydrophilic

polyoxyethylene (POE). This results in an octablock copolymer with the following general formula:



wherein:

a is a number such that the hydrophile portion represented by polyoxyethylene $(C_2H_4O)_a$ (POE) constitutes between approximately 10% to 40% of the total molecular weight of the compound;

the mean aggregate molecular weight of the hydrophobe portion of the octablock copolymer consisting of polyoxypropylene $(C_3H_6O)_b$ (POP) is between approximately 5000 and 7000 daltons; and

b is a number such that the polyoxypropylene $(C_3H_6O)_b$ (POP) portion of the total molecular weight of the octablock copolymer constitutes between approximately 60% and 90% of the compound.

A particularly useful composition is an admixture of a compound capable of altering gene expression and/or protein translation, such as an antisense oligonucleotide, a triplex DNA compound, a ribozyme or other compound capable of altering nucleic acid sequence function, and either of the above-described nonionic block copolymer, the above-described POE/POP copolymer, or a combination thereof.

The composition of the present invention can be administered by a number of routes including, but not limited to topical, transdermal, oral, trans-mucosal, subcutaneous injection, intravenous injection, intraperitoneal injection and intramuscular injection.

Accordingly, an object of the invention is to provide a therapeutic drug delivery vehicle.

Another object of the present invention is to provide compositions that facilitate delivery of one or more therapeutic nucleic acid sequence function altering agents into the interior of a cell, such as a phagocytic cell, when admixed with a therapeutic agent.

Another object of the present invention is to provide compositions that act synergistically with a delivered agent once inside a cell.

Still another object of the invention is to provide nonionic block copolymers and POE/POP copolymers having surfactant properties that facilitate the transmission and introduction across cellular plasma membranes of nucleic acid sequences and compounds capable of altering nucleic acid sequence function.

A further object of the present invention is to provide compositions and a method for treating genetic and physiologic disorders using nucleic acid sequences and antisense oligonucleotides in combination with nonionic block copolymers, POE/POP copolymers, or a combination thereof.

Another object of the present invention is to provide compositions and a method useful for manipulating the expression of genes using triplex DNA compounds.

Yet another object of the invention is to provide DNA vaccines.

It is an object of the present invention to provide compositions which can be used to treat persons with infectious diseases.

Yet another object of the present invention is to provide a method of treating viral infections in humans or animals.

Another object of the present invention is a compound and method that is effective in inhibiting the replication of viruses in both animals and humans.

Another object of the present invention is to provide a compound and method that is effective in inhibiting the replication of HIV and other RNA and DNA viruses.

5 Yet another object of the present invention is to provide a method of treating microbial infections in humans or animals.

It is another object of the present invention to inactivate virus in a blood product prior to infusion into a person or animal.

10 These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiment and the appended claims.

15 **Brief Description of the Drawings**

Fig. 1 is a grid illustrating block copolymers by molecular weight of hydrophobe and percent hydrophile.

20 Fig. 2 is a grid illustrating preferred therapeutic delivery block copolymers by molecular weight of hydrophobe and percent hydrophile.

Fig. 3 is a grid illustrating more preferred therapeutic delivery block copolymers by molecular weight of hydrophobe and percent hydrophile.

25 Fig. 4 picture of results of experiments with administration of nucleic acids and polymers.

Fig. 5 is a graph of transfection results

Detailed Description

30 The present invention includes gene therapy compositions that are admixtures of a nonionic block copolymer, a POE/POP copolymer, or a combination thereof, and nucleic acid sequences or compounds capable of altering nucleic acid sequence function, and methods of delivering these compositions to a human or animal in need thereof for the

intracellular alteration of gene expression and/or protein translation.

It has been unexpectedly found that high molecular weight surface active nonionic polyoxyethylene-polyoxypropylene block copolymers having a low percentage of polyoxyethylene, POE/POP copolymers, and a combinations thereof, facilitate the transport of DNA and other compounds into cells and thus are useful for the intracellular delivery of therapeutic agents *in vivo* for the treatment of disease. It is believed that both the block copolymers and the POE/POP copolymers are particularly useful in helping to reseal membranes and thus increase the percent survival of cells wherein nucleic acid sequences or other compounds have been intracellularly introduced. Surprisingly, it has also been found that compositions comprising the nonionic block copolymers and/or POE/POP copolymers of the present invention and nucleic acid sequences are less susceptible to the degrading effects of DNAase than nucleic acid sequences alone.

The present invention also comprises therapeutic compositions and methods which kill or inhibit the growth of microorganisms and alter the expression or function nucleic acid sequences. An example of the bacteria that the present invention is effective against is mycobacteria species, such as *Mycobacterium tuberculosis*, *Mycobacterium avium*, and *Mycobacterium leprae*. Other microorganisms that the invention is effective against include, but are not limited to, *Chlamydia trachomatis*, *Chlamydia pneumoniae*, *Listeria monocytogenes*, *Candida albicans*, *Cryptococcus neoformans*, *Toxoplasma gondii*, *Pneumocystis carinii*, *Herpes simplex virus* type 1, *Cytomegalovirus*, influenza virus type A and B, and respiratory syncytial virus.

The present invention includes therapeutic compositions and methods for treating DNA viruses and RNA viruses, and infections and infectious diseases caused by such viruses in a human or animal, including infections caused by

HIV or herpes or antigenically-related strains thereof. Antigenically-related strains are strains that cross react with antibodies specific for HIV. One skilled in the art can readily determine viral strains that are antigenically-related to HIV by
5 conducting standard immunoassay tests using anti-HIV antibodies and the viral strain to be analyzed, and looking for positive cross-reactivity. The therapeutic compositions comprising the surface active copolymers and/or POE/POP copolymers disclosed herein are effective in inhibiting or
10 suppressing the replication of such viruses in cells.

The present invention includes a therapeutic composition useful for delivering antimicrobial drugs and treating disease states comprising an admixture of a surface active nonionic block copolymer, a POE/POP copolymer, or a
15 combination thereof; a compound capable of altering nucleic acid sequence function; and an antibiotic or therapeutic drug. Examples of such compounds capable of altering nucleic acid sequence function include genes, oligonucleotides, antisense oligonucleotides, triplex DNA compounds, and ribozymes.
20 Drugs that can be used with the nonionic copolymers of the present invention include, but are not limited to, rifampin, isoniazid, ethambutol, gentamicin, tetracycline, erythromycin, pyrazinamide, streptomycin, clofazimine, rifabutin, fluoroquinolones such as ofloxacin and sparfloxacin,
25 azithromycin, clarithromycin, dapsone, doxycycline, ciprofloxacin, ampicillin, amphotericin B, fluconazole, ketoconazole, fluconazole, pyrimethamine, sulfadiazine, clindamycin, azithromycin, paromycin, diclazaril, clarithromycin, atovaquone, pentamidine, acyclovir,
30 trifluorouridine, AZT, DDI, DDC, and other antiviral nucleoside analogs, foscarnat, ganciclovir, viral protease inhibitors, antisense and other modified oligonucleotides, and ribavirin.

Preferred drugs to use for various infectious
35 microorganisms are listed in Table I.

Table I

	<u>Organism</u>	<u>Drugs</u>
5	<u><i>Bacteria</i></u>	
10	Mycobacterium tuberculosis	Isoniazid, rifampin, ethambutol, pyrazinamide, streptomycin, clofazimine, rifabutin, fluoroquinolones such as ofloxacin and sparfloxacin

2025-04-24 14:44:44

Table I (continued)

	<u>Organism</u>	<u>Drugs</u>
5	<i>Mycobacterium avium</i>	Rifabutin, rifampin, azithromycin, clarithromycin, fluoroquinolones
	<i>Mycobacterium leprae</i>	Dapsone
10	<i>Chlamydia trachomatis</i>	Tetracycline, doxycycline, erythromycin, ciprofloxacin
	<i>Chlamydia pneumoniae</i>	Doxycycline, erythromycin
15	<i>Listeria monocytogenes</i>	Ampicillin
	<u>Fungi</u>	
20	<i>Candida albicans</i>	Amphotericin B, ketoconazole, fluconazole
	<i>Cryptococcus neoformans</i>	Amphotericin B, ketoconazole, fluconazole
25	<u>Protozoa</u>	
	<i>Toxoplasma gondii</i>	Pyrimethamine, sulfadiazine, clindamycin, azithromycin, clarithromycin, atovaquone
30	<i>Pneumocystis carinii</i>	Pentamidine, atovaquone
35	<i>Cryptosporidium</i> sp.	Paromomycin, diclazaril
	<u>Virus</u>	
40	Herpes simplex virus type 1 and type 2	Acyclovir, trifluorouridine and other antiviral nucleoside analogs, foscarnat, antisense oligonucleotides, and triplex-specific DNA sequences
45	Cytomegalovirus	Foscarnet, ganciclovir
	HIV	AZT, DDI, DDC, foscarnat, viral protease inhibitors, peptides, antisense oligonucleotides, triplex and other nucleic acid sequences
50		

Table I (continued)

	<u>Organism</u>	<u>Drugs</u>
5	Influenza virus type A and B	Ribavirin
	Respiratory syncytial virus	Ribavirin
10	Varizella zoster virus	Acyclovir


15 Optionally, surfactants and low molecular weight alcohols are added to the therapeutic admixture of antimicrobial drug and nonionic block copolymer, POE/POP copolymer, or combination thereof. Examples of surfactants useful in the present invention include Tween 80 and emulsions with fatty acids such as phospholipids, cholate and amino acids. The preferred surfactant is Tween 80. Surfactants are added to the admixture at a concentration ranging from approximately 0.1% to approximately 5% v/v. The preferred surfactant concentration is approximately 2%. The term "approximately" as it applies to concentrations expressed herein means the stated concentration plus or minus ten percent. The term "low molecular weight alcohols" means alcohols having two to eight carbons. An example of a low molecular weight alcohols useful in the present invention is ethanol, which is the preferred low molecular weight alcohol. Low molecular weight alcohols are added to the admixture at a concentration ranging from approximately 0.5% to approximately 5% v/v. The preferred low molecular weight alcohol concentration is between approximately 1% and approximately 3% v/v.

35 The present invention also includes compositions and methods for immunizing animals or humans, otherwise termed DNA vaccination. Immunization is accomplished by administering a composition comprising the gene that codes for the gene product to be immunized against contained in an expression, in combination with a block copolymer that

promotes and facilitates uptake of genetic material across cell membranes. The introduced gene is expressed, resulting in the production of an antigenic gene product.

5 Still further, compositions comprising nonionic block copolymers, POE/POP copolymers, or a combination thereof and genes that code for compounds effective for killing, reducing or retarding cancer, such as lymphokines, may be administered to humans or animals for the treatment of cancer.


10 The present invention comprises a surface active copolymer that is preferably an ethylene oxide-propylene oxide condensation product with the following general formula:



15 wherein a is an integer such that the hydrophobe represented by (C₃H₆O) has a molecular weight of between approximately 750 and approximately 15,000, and b is an integer such that the hydrophile portion represented by (C₂H₄O) constitutes approximately 1% to approximately 50% by weight of the compound.

20 The present invention also comprises a therapeutic delivery composition useful for altering gene expression and/or protein translation comprising an administerable admixture of an effective amount of an anti-sense oligonucleotide or other nucleic acid sequence, and an effective amount of a nonionic block copolymer having the following general formula:


25



30 wherein a is an integer such that the hydrophobe represented by (C₃H₆O) has a molecular weight of approximately 750 and approximately 15,000, preferably between approximately 2250 and approximately 15,000, more preferably between approximately 3250 and approximately 15,000, and b is an integer such that the hydrophile portion represented by (C₂H₄O) constitutes approximately 1% to approximately 50% by weight


of the compound, preferably approximately 5% to approximately 20%. The term admixture as used herein means any combination of therapeutic drug and nonionic block copolymer, including solutions, suspensions, or encapsulations of drug in copolymer micelles. An effective amount is an amount sufficient to alter the activity and/or the amount of gene product produced by the gene or genes sought to be modulated in a human or animal.

The present invention also comprises a therapeutic delivery composition useful for immunizing an animal or human against a particular gene product comprising an administerable admixture of an effective amount of an expression vector, the gene that codes for the gene product to be immunized against contained in the expression vector, and an effective amount of a nonionic block copolymer having the following general formula:



wherein a is an integer such that the hydrophobe represented by $(\text{C}_3\text{H}_6\text{O})$ has a molecular weight of approximately 750 and approximately 15,000, preferably between approximately 2250 and approximately 15,000, more preferably between approximately 3250 and approximately 15,000, and b is an integer such that the hydrophile portion represented by $(\text{C}_2\text{H}_4\text{O})$ constitutes approximately 1% to approximately 50% by weight of the compound, preferably approximately 5% to approximately 20%. An effective amount is an amount sufficient to elicit an immunological response against the gene product of the nucleic acid sequence administered to the human or animal.

The surface active nonionic block copolymer of the present invention comprises a reverse triblock copolymer of polyoxypropylene/polyoxyethylene having the formula:



POP POE POP

wherein "b" represents a number such that the molecular weight of the hydrophobe (C₃H₆O)_b is between approximately 2,000 and 10,000, and "a" represents a number such that the percentage of hydrophile (C₂H₄O)_a is between approximately 2% and 30%.


Another embodiment of the surface active nonionic block copolymer present invention, comprises the reverse triblock copolymer of polyoxypropylene/polyoxyethylene having the formula:



POP POE POP

wherein "b" represents a number such that the molecular weight of the hydrophobe (C₃H₆O)_b is between approximately 2,500 and 8,500, and "a" represents a number such that the percentage of hydrophile (C₂H₄O)_a is between approximately 7% and 23%.


Yet another embodiment of the present invention, comprises the reverse triblock copolymer of polyoxypropylene/polyoxyethylene having the formula:



POP POE POP

wherein "b" represents a number such that the molecular weight of the hydrophobe (C₃H₆O)_b is between approximately 3,000 and 7,000, and "a" represents a number such that the percentage of hydrophile (C₂H₄O)_a is between approximately 8% and 20%.

A preferred embodiment of the present invention, comprises the reverse triblock copolymer polyoxypropylene/polyoxyethylene having the formula:



POP POE POP

wherein "b" represents a number such that the molecular weight of the hydrophobe $(C_3H_6O)_b$ is between approximately 5,000 and 6,000, and "a" represents a number such that the percentage of hydrophile $(C_2H_4O)_a$ is between approximately 14% and 16%.

In general, "a" represents a number such that the percentage of hydrophile $(C_2H_4O)_a$ is between approximately 5% and 30%, with a preferable range of between approximately 7% and 23% and a most preferable range of between 8% and 20%.

It should be understood that the molecular weight and percentage ranges that are described for the block copolymer are to be considered outside ranges and that any population of molecules that falls within the stated ranges is considered an embodiment of the present invention.

The entire block copolymer molecule is poorly soluble in water and is substantially nonionic. The steric configurations and physiochemical properties of the molecule, rather than the chemical nature of the constituent parts, are believed to be largely responsible for the antiinfective activity and therapeutic delivery activity. Compositions of the present invention include, but are not limited to aqueous solutions, suspensions or emulsions, such as oil-in-water emulsions.

The polymer blocks are formed by condensation of ethylene oxide and propylene oxide, at elevated temperature and pressure, in the presence of a catalyst. There is some statistical variation in the number of monomer units which combine to form a polymer chain in each copolymer. The molecular weights given are approximations of the average weight of copolymer molecule in each preparation and are dependent on the assay methodology and calibration standards used. It is to be understood that the blocks of propylene oxide and ethylene oxide do not have to be pure. Small amounts of other materials can be admixed so long as the overall physical chemical properties are not substantially changed. A more

detailed discussion of the preparation of these products is found in U.S. Patent No. 2,674,619, which is incorporated herein by reference in its entirety.

5 Ethylene oxide-propylene oxide condensation
products which may be employed in the present invention are
summarized in Table II. It is to be understood that these
compounds are merely representative of the compounds that can
be used to practice the present invention and do not include all
possible compounds that could be used to practice the present
10 invention. The high molecular weight copolymers listed in
Table II that do not have a BASF tradename are novel
compositions that have never been synthesized before.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
22

Table II

CRL #	Poloxamer	BASF Trade Name	Molecular Weight of POP	% POE
	122	L42	≈1200	≈20%
CRL-85171	181	L61	≈1750	≈10%
CRL-85172	182	L62	≈1750	≈20%
CRL-85173	183	L63	≈1750	≈30%
CRL-85174	184	L64	≈1750	≈40%
CRL-85175	185	P65	≈1750	≈50%
CRL-85178	188	F68	≈1750	≈80%
CRL-85202	212	L72	≈2050	≈20%
CRL-85221	231	L81	≈2250	≈10%
CRL-8122	282	L92	≈2750	≈20%
CRL-8131	331	L101	≈3250	≈10%
CRL-8133	333	P103	≈3250	≈30%
CRL-8135	335	P105	≈3250	≈50%
CRL-9038	338	F108	≈3250	≈80%
CRL-8141	401	L121	≈4000	≈10%
CRL-8142	402	L122	≈4000	≈20%
CRL-8143	403	P123	≈4000	≈30%
CRL-8941	441	L141	≈4400	≈10%
CRL-8950	-----	-----	≈6000	≈5%
CRL-1235	-----	-----	≈7500	≈5%
CRL-1190	-----	-----	≈10,000	≈5%
CRL-336	-----	-----	≈14,000	≈5%
CRL-1183	-----	-----	≈3750	≈10%
CRL-1122	-----	-----	≈5900	≈12%
CRL-3362	-----	-----	≈3900	≈11%
CRL-3632	-----	-----	≈4740	≈11%
CRL-9352	-----	-----	≈7750	≈15%
CRL-1187	-----	-----	≈750	≈25%

5 A grid illustrating the range of copolymer
encompassed by the present invention based upon the molecular
weight of the hydrophobe portion and the percent hydrophile,
and showing selected nonionic block copolymers appears as
Figure 1. The polymer blocks are formed by condensation, at
elevated temperature and pressure, of ethylene oxide and
propylene oxide in the presence of a catalyst. There is some
statistical variation in the number of monomer units which
10 combine to form a polymer chain in each copolymer. The
molecular weights given are approximations of the average size
of copolymer molecules in each preparation. A further
description of the preparation of these block copolymers is
found in U.S. Patent No. 2,674,619. (Also see, "A Review of
15 Block Polymer Surfactants", Schmolka I.R., *J. Am. Oil Chemist
Soc.*, 54:110-116 (1977) and *Block and Graft Copolymerization*,
Volume 2, edited by R.J. Ceresa, John Wiley and Sons, New
York, 1976.


20 It has been discovered that the copolymers
particularly effective as therapeutic delivery agents are shown
in Figures 2 and 3. As is apparent from Figures 2 and 3, the
copolymers most effective as therapeutic delivery agents are
high molecular weight and have low percentages of POE -
generally less than 20% POE.

25 Non-ionic block copolymers form micelles above
their critical micelle concentration. The non-ionic copolymers
have negative thermal coefficients of solubility. In the cold, the
kinetic energy of water molecules is reduced and they form
weak hydrogen bonds with the oxygen of the POP blocks. This
30 hydration of the hydrophobe promotes solubility at low
temperatures. As the temperature rises, the "cloud point" is
reached; the increased kinetic energy of the water breaks the
hydrogen bonds, the polymer becomes insoluble and micelles
form.


The biologically-active POE/POP copolymers of the present invention comprise a surface active compound with four hydrophobic segments and a small proportion of hydrophile. Typical examples have eight segments or octablock structure with a core of either a hydrophobic or hydrophilic central structure and a hydrophilic or hydrophobic outer structure.

The entire molecule is poorly soluble in water and is either a nonionic or weakly cationic surface active agent. The steric configuration and physiochemical properties of the molecule, rather than the chemical nature of the constituent parts, are thought to be responsible for the biologic effects of the copolymer.

The POE/POP copolymers of the present invention comprise blocks of polyoxypropylene and polyoxyethylene built on an alkylenediamine initiator. The blocks of polyoxypropylene (POP) and polyoxyethylene (POE) have the following structures:



POLYOXYPROPYLENE (POP)




POLYOXYETHYLENE (POE)

The polymer blocks are formed by condensation of ethylene oxide and propylene oxide onto a tetrafunctional ethylene diamine initiator at elevated temperature and pressure in the presence of a basic catalyst. There is some statistical

variation in the number of monomer units which combine to form a polymer chain in each copolymer. The molecular weights given are approximations of the average weight of copolymer molecule in each preparation. A further description of the preparation of these block copolymers is found in U.S. Patent No. 2,674,619 and U.S. Patent No. 2,979,528. (Also see "A Review of Block Polymer Surfactants", Schmolka, I.R., *J. Am. Oil Chemists' Soc.*, 54:110-116 (1977) and *Block and Graft Copolymerization*, Volume 2 edited by R.J. Ceresa, John Wiley & Sons, New York (1976).

In one aspect of the biologically active POE/POP copolymers of the present invention, the block copolymer comprises a polymer of hydrophobic polyoxypropylene (POP) built on an ethylenediamine initiator. Polymers of hydrophilic polyoxyethylene (POE) are then built on the block of hydrophobic polyoxypropylene (POP). This results in an octablock copolymer with the following general formula:



wherein:


the mean aggregate molecular weight of the hydrophobe portion of the octablock copolymer consisting of polyoxypropylene (C₃H₆O)_b (POP) is between approximately 5000 and 7000 daltons;

a is a number such that the hydrophile portion represented by polyoxyethylene (C₂H₄O)_a (POE) constitutes

between approximately 10% to 40% of the total molecular weight of the compound; and

b is a number such that the polyoxypropylene (C₃H₆O)_b (POP) portion of the total molecular weight of the octablock copolymer constitutes between approximately 60% and 90% of the compound.

In another aspect of the present invention, the POE/POP block copolymer comprises a polymer of hydrophilic polyoxyethylene (POE) built on an ethylene diamine initiator. Polymers of hydrophobic polyoxypropylene (POP) are then built on the block of hydrophilic polyethylene (POE). This results in an octablock copolymer with the following general formula:



wherein:

The molecular weight of the hydrophobe portion of the octablock copolymer consisting of polyoxypropylene (C₃H₆O)_b (POP) is between approximately 5000 and 7000 daltons;

a is a number such that the hydrophile portion represented by polyoxyethylene (C₂H₄O)_a (POE) constitutes between approximately 10% and 40% of the total molecular weight of the compound; and

b is a number such that the polyoxypropylene (C₃H₆O)_b (POP) portion of the octablock copolymer constitutes between approximately 60% and 90% of the compound.

This type of polymer is called reverse copolymer because its structure is the reverse of octablock copolymers that have polyoxypropylene (POP) in the center flanked by blocks of polyoxyethylene (POE).


5 The (C_3H_6O) portion of the copolymer can constitute up to 95% of the octablock copolymer. The (C_2H_4O) portion of the copolymer can constitute as low as 5% of the Octablock copolymer.

10 The octablock copolymers comprising the biologically active copolymers of the present invention include, but are not limited to, the block copolymers Tetronic® and reverse Tetronic® manufactured by the BASF Corporation (BASF Corporation, Parsippany, NJ). These include copolymers wherein:

15 a is a number such that the hydrophile portion represented by polyoxyethylene $(C_2H_4O)_a$ (POE) constitutes between approximately 5% and 20% of the total molecular weight of the compound;

20 b is a number such that the polyoxypropylene $(C_3H_6O)_b$ (POP) portion of the octablock copolymer constitutes between approximately 80% and 95% of the compound.

25 A preferred biologically active copolymer is the octablock copolymer T110R1 (BASF Corporation, Parsippany, NJ) which corresponds to the following formula:



30


wherein:

The mean molecular weight of the hydrophobe portion of the octablock copolymer represented by polyoxypropylene (C₃H₆O)_b (POP) is approximately 5,220 daltons;

5 a is a number such that the hydrophile portion represented by polyoxyethylene (C₂H₄O)_a (POE) constitutes approximately 10% of the compound by weight; and

10 b is a number such that the polyoxypropylene (C₃H₆O)_b (POP) portion of the octablock copolymer constitutes approximately 90% of the compound by weight.

A preferred biologically active copolymers is the octablock copolymer T130R2 (BASF Corporation, Parsippany, NJ) which corresponds to the following formula:



wherein:


20 The mean molecular weight of the hydrophobe portion of the octablock copolymer represented by polyoxypropylene (C₃H₆O)_b (POP) is approximately 5750 daltons;

25 a is a number such that the hydrophile portion represented by polyoxyethylene (C₂H₄O)_a (POE) constitutes approximately 20% of the compound by weight; and

 b is a number such that the polyoxypropylene (C₃H₆O)_b (POP) portion of the octablock copolymer constitutes approximately 80% of the compound by weight.

30 Another preferred embodiment of the biologically active copolymers of the present invention is the compound

designated T1501 (BASF Corporation, Parsippany, NJ) which corresponds to the following formula:




wherein:

the mean molecular weight of the hydrophobe portion of the octablock copolymer represented by polyoxypropylene (C₃H₆O)_b (POP) is approximately 6750 daltons;

a is a number such that the hydrophile portion represented by polyoxyethylene (C₂H₄O)_a (POE) constitutes approximately 10% of the compound by weight; and

b is a number such that the polyoxypropylene (C₃H₆O)_b (POP) portion of the octablock copolymer constitutes approximately 90% of the compound by weight.

The most preferred embodiment of the biologically active copolymers of the present invention is the octablock copolymer T150R1 (BASF Corporation, Parsippany, NJ) which corresponds to the following formula:



wherein:

The mean molecular weight of the hydrophobe portion of the octablock copolymer represented by polyoxypropylene (C₃H₆O)_b (POP) is approximately 6750 daltons;

a is a number such that the hydrophile portion represented by polyoxyethylene (C₂H₄O)_a (POE) constitutes approximately 10% of the compound by weight; and

b is a number such that the polyoxypropylene (C₃H₆O)_b (POP) portion of the octablock copolymer constitutes approximately 90% of the compound by weight.

The present invention also comprises a therapeutic delivery composition useful for immunizing an animal or human against a particular gene product comprising an administerable admixture of an effective amount of an expression vector, the gene that codes for the gene product to be immunized against contained in the expression vector, and an effective amount of a POE/POP copolymer of the present invention.

The present invention also comprises a therapeutic delivery composition useful for immunizing an animal or human against a particular gene product comprising an administerable admixture of an effective amount of an expression vector, the gene that codes for the gene product to be immunized against contained in the expression vector, and an effective amount of a POE/POP copolymer of the present invention.

The POE/POP copolymers of the present invention, including biological activity thereof, are further discussed in U.S. Patent No. 5,494,660, which is incorporated herein in its entirety by reference.

Thus, the copolymers, including both the nonionic block copolymers and the POE/POP copolymers of the present invention, which are therapeutic themselves, can form physical structures that can be combined or loaded with an additional,

distinct therapeutic agent. Consequently, the nonionic block copolymers and the POE/POP copolymers of the present invention can be used as therapeutic drug delivery vehicles. Admixtures of therapeutic drugs with non-ionic block and POE/POP copolymers have the advantage of synergistic activity of at least two therapeutic agents. Further, copolymers having specific characteristics can be selected for use with particular therapeutic drugs. For example, CRL-8131, which is hydrophobic, is an excellent carrier for hydrophobic antibiotics such as rifampin. However, other agents which are not distinctly hydrophobic can be used according to the present invention.

A therapeutic delivery vehicle is prepared using any of the surface active nonionic block copolymers and/or POE/POP copolymers of the present invention in combination with any of a variety of antimicrobial agents. In a one embodiment CRL-8131 is used at a concentration of approximately 3% to approximately 5% to construct a therapeutic delivery vehicle. Therapeutic delivery vehicles made using copolymers that are more hydrophilic than CRL-8131 normally require a higher concentration (approximately 5% to approximately 10%) of the copolymer.

Using copolymer-based micelles as a therapeutic drug delivery vehicle is particularly desirable because the micelles are accumulated readily and are present for an extended period of time, in macrophages, the site of HIV and other viral infections and a major target for viral therapy. Examples of such therapeutic copolymer-based therapeutic compositions include CRL-8131 combined with 2% Tween 80 and 1% ethanol, and CRL-8142 combined with 1% Tween 80 and 5% ethanol.

Nucleic acid sequences or other compounds capable of altering nucleic acid sequence function are administered to a human or animal to alter gene expression and/or modify the amount or activity of gene product. For

example, antisense oligonucleotides admixed with either or both of the above-described copolymers yield compositions useful for delivery of the antisense oligonucleotides for the purpose of altering or regulating gene expression and/or protein translation. Further, nucleic acid sequences such as genes can be administered which incorporate into the chromosome replacing or augmenting the defective gene. Alternatively, the intracellularly administered gene may reside in the cell and be expressed in an extrachromosomal element.

The present invention also provides novel compositions and methods for immunizing an animal or human. The compositions comprise an expression vector, a gene that codes for the gene product to be immunized against contained in the expression vector, and a block copolymer, which includes the nonionic block copolymers, the POE/POP copolymers, or a combination thereof, effective for transferring genetic material such as expression vectors across the membrane of cells. The method of immunizing an animal or human comprises administering of the expression vector-containing copolymer composition to the animal or human. A preferred mode of administration is by intraperitoneal injection. This embodiment of the invention provides means for the delivery of genetic sequences capable of expressing an antigenic gene product directly into human or animal cells, either *in vivo* or *ex vivo* with subsequent reintroduction into the human or animal. Once introduced into the cells the production of antigenic gene product induces and maintains an immune response by the human or animal against the introduced gene product.

The following specific examples illustrate various aspects of the invention, such as compositions and methods of the invention useful for gene therapy, and compositions and methods of the invention useful for gene-mediated immunization. It should be appreciated that other embodiments and uses will be apparent to those skilled in the art and that the invention is not limited to these specific illustrative examples.

Example I

5 A therapeutic delivery vehicle is prepared by combining any of the surface active nonionic block copolymers, such as CRL-8131 with any of a variety of compounds capable of altering nucleic acid sequence function. For CRL-8131 a concentration of three to five percent weight per volume is desirable to construct the therapeutic vehicle. For more hydrophilic copolymer a five to ten percent weight per volume.

10 300 milligrams of CRL-8131 was added to 10 ml of 0.9% NaCl and the mixture is solubilized by storage at temperatures of 2-4°C until a clear solution is formed. An appropriate amount of a compound capable of altering nucleic acid gene function is added to the mixture and micelles associating the copolymer and the compound are formed by raising the temperature above 5°C and allowing the suspension of micelles to equilibrate. The equilibrated suspension is suitable for administration.

15 For example, an antisense oligonucleotide sequence, such as one of those disclosed by Matsukara, M. et al. *Proc. Natl. Acad. Sci. USA* 84:7706-7710 (1987), which is expressly incorporated herein in its entirety by reference, is combined with the copolymer to form a micelle composition.

20 Briefly, phosphorothioate or methylphosphonate derivatives of a sequence complimentary to regions of the *art/trs* genes of HIV having the sequence 5'-TCGTCGCTGTCTCG-3' are prepared according to the method of Matsukura et al. 300 milligrams of CRL-8131 is added to 10 ml of 0.9% NaCl and the mixture is solubilized by storage at temperatures of 2-4°C until a clear solution is formed. The desired antisense oligonucleotide subsequently is mixed with the copolymer solution to provide a concentration effective in inhibiting viral activity when administered to a patient infected with the HIV virus. Generally the effective amount of antisense compound will be such that the final concentration in the blood

25
30
35

is in the range of 1 μ M to 100 μ M, although other effective amounts of antisense compounds outside this range may be found for specific antisense compounds. One skilled in the art can readily test the relative effectiveness of any particular antisense oligonucleotide according to the *in vivo* test of Matsukura, et al.

An average person has approximately 6.25 liters of blood. Thus, oligonucleotide concentrations of approximately 6 mM to 600 mM are required in the composition when 1 ml injections are to be administered. Lower oligonucleotide compositions can be used with larger administration volumes.

Example II

The antiinfective antisense oligonucleotide composition of Example I is administered to HIV patients by any route effective to reduce viral activity. The preferred route of administration is by intravenous injection. The antisense composition may be administered multiple times a day to ensure that an effective amount of the antisense oligonucleotide is maintained.

Example III

A gene therapy composition for treating an animal or human suffering from the effects of a defective or missing gene is made by combining a copolymer, such as CRL-8131 with a normal copy of the defective gene. For example, for patients suffering from adenosine deaminase (ADA) deficiency a gene therapy composition is made that contains a normal copy of the adenosine deaminase gene. The gene therapy composition is made by mixing a copolymer prepared as described above in Example I with the desired gene, removing blood from the human or animal, transfecting blood cells with the ADA gene-containing composition, and reintroducing the transfected blood cells into the human or animal. The

introduced gene is expressed *in vivo*, alleviating the effects of the original gene deficiency.

Example IV

5 Similarly, the gene therapy composition of Example III is combined with isolated T-lymphocytes to form T-lymphocytes containing the ADA gene. The ADA gene-containing T-lymphocytes are subsequently administered, for example by injection, into the patient suffering from adenosine
10 deaminase deficiency. The administered cells express the ADA and produce adenosine deaminase, thus augmenting the supply of the enzyme in the patient and correcting the deficiency.

Example V

15 DNA vaccination is carried out essentially as described for gene therapy in Examples III or IV, except that the gene that is introduced into the host expresses an antigenic gene product that is recognized as foreign by the host animal, thus eliciting an immune response.

Example VI

 A composition comprising copolymer CRL-8131 and an expression vector containing the gD gene of *Herpes simplex* virus type-1 was used in a transfection experiment.
25 DNA transfection normally is performed using standard calcium chloride and DEAE dextran precipitation techniques. DEAE dextran is used to rough up the cell membrane and calcium is used to precipitate DNA onto the cell surface, facilitating DNA uptake into the cells. This procedure is
30 generally toxic to the cells, however, and causes substantial cellular mortality.

 A new transfection system was discovered using the block copolymer of the present invention in place of calcium chloride. In fact, it was surprisingly discovered that

copolymer assisted transfection occurs even in the absence of DEAE dextran.

5 Vero cells were incubated in DEAE dextran for 30 seconds. A mixture of copolymer and an expression vector containing glycoprotein gD DNA of *Herpes simplex* virus type-1 was added to Vero cells immediately after the removal of DEAE dextran. It was found that up to 40% of the cells were effectively transfected with the gD gene.

10 Surprisingly, in two out of four experiments copolymers were able to transfect Vero cells at a lower than 40% efficiency even when the DEAE dextran incubation step is omitted.

Example VII

15 Other studies have also demonstrated that block copolymers are effective in transferring genetic material across cellular membranes *in vivo*. DNA vaccine-induced immunization was successful when an expression vector containing the gD gene of *Herpes simplex* virus type-1 combined with copolymer was injected intraperitoneally into rabbits every two weeks. Sera was collected and tested for the presence of anti-gD antibody. Low levels of anti-gD antibody were detected after 4 weeks of inoculation in this fashion. These results demonstrate that genetic material administered intraperitoneally with a copolymer delivery vehicle is taken up by cells *in vivo* and expressed to give the gene product in quantities sufficient to elicit an immune response.

Example VIII

30 **DNase protection experiments.** Five different compounds (CRL 1122, 3362, 3632, 9352, and 8131) were used in experiments to test the degree of protection. DNA was mixed with compounds at 4°C, and after 15 min. at 37°C DNase I (1 µl of 10 mg/ml solution) was added. After 30 min. of incubation at 37°C, DNase I was removed by treatment with

35

proteinase K (3 μ l of 10 mg/ml solution). Controls were: DNase I in the absence of nonionic block copolymer and DNA alone without any DNase I treatment.

5 DNA was protected from DNase I degradation in all samples in which nonionic block copolymers were present. The best protection of DNA was achieved with CRL-3362 and 8131. DNA copolymer compositions did not migrate in horizontal agarose electrophoresis and remained within the wells (stained with ethidium bromide). Effective protection
10 against DNase I action was achieved in solutions of 1 volume DNA solution (1 μ g/ml) to 5 volumes of nonionic block copolymer (30 μ g/ml). The estimated amount of protection varied from experiment to experiment and was estimated to be within 15-40% of total DNA.

15 Additional experiments showed that DNA-copolymer compounds failed to transform *E. coli* competent cells via the calcium method. Phenol also failed to dissolve nonionic block copolymer away from DNA. DNA bound to NBC can be precipitated by adding 5 volumes of isopropyl
20 alcohol.

Example IX

Transfection experiments. Typical transfection experiments for transient expression of herpes viral glycoprotein genes and other genes of interest involved the
25 following procedure. Cells such as COS (African monkey kidney cells; CV1) are seeded on 6-well plates. Transfection is performed when cells are 50-80% confluent (still in log growth phase). Cells are first washed with PBS buffer, they are
30 incubated with 0.5 ml of DEAE-Dextran solution (500 mg/ml) for 1-2 minutes, this solution is aspirated and DNA precipitate is added to cells. DNA to be transfected is mixed for 30 min. at room temperature with CaCl_2 at controlled pH conditions to form a fine precipitate. This solution is mixed with 1 ml of
35 growth medium (DMEM) and put onto cells for 4 hours at

37°C. At this time, the cells are shocked with 15% glycerol and subsequently washed with PBS. This osmotic shock facilitates the taking up of CaCl_2 -DNA precipitate into cells. Cells are then washed again with PBS, and incubated with growth media at 37°C for 48 hours.

Gene expression is detected in most cases using specific monoclonal antibodies directed against the expressed proteins using indirect immunofluorescence. The expressed proteins can be also labeled with radioactive tracers and immunoprecipitated or detected in westerns.

25 µl of DNA (7 µg) and 25 µl of nonionic block copolymer (30 µg/ml) were used. Additionally, mixing of nonionic block copolymer with DNA on ice, and addition of mixture into the cells produced similar results as when they were added separately (DNA added first and nonionic block copolymer second).

Copolymers 1183, 1187, 8131, 1235, 8950AQ and 1190AQ (where AQ indicates that the nonionic block copolymers were diluted 1:10 and 25 µl were used). Typical results are as follows. Transfection with DNA alone, dextran alone, copolymer alone, and DNA plus dextran had negligible transfection of less than 0.2%. In contrast, the positive control of DNA plus dextran plus glycerol has transfection of 2% while various copolymers plus DNA were successful in transfecting DNA into cells up to 2.5 times better than the control, as shown in Table III:

Table III

Copolymer	Percent Transfection
1183	2%
1187	5%
8131	2%
1235	3%
8950AQ	4%
1190AQ	5%

5 There was no copolymer associated toxicity
except mild toxicity with 1187. The others were toxic
especially after glycerol treatment.

10 It should be understood that the foregoing relates
only to preferred embodiments of the present invention and that
numerous modifications and alterations may be made without
departing from the spirit and scope of the invention as set forth
in the appended claims.